

## Table Of Content

|   |   |
|---|---|
| <b>Journal Cover</b> .....                  | 2 |
| <b>Author[s] Statement</b> .....            | 3 |
| <b>Editorial Team</b> .....                 | 4 |
| <b>Article information</b> .....            | 5 |
| Check this article update (crossmark) ..... | 5 |
| Check this article impact .....             | 5 |
| Cite this article .....                     | 5 |
| <b>Title page</b> .....                     | 6 |
| Article Title .....                         | 6 |
| Author information .....                    | 6 |
| Abstract .....                              | 6 |
| <b>Article content</b> .....                | 7 |

**ISSN (ONLINE) 2598-9936**



**INDONESIAN JOURNAL OF INNOVATION STUDIES**  
PUBLISHED BY  
UNIVERSITAS MUHAMMADIYAH SIDOARJO

## Originality Statement

The author[s] declare that this article is their own work and to the best of their knowledge it contains no materials previously published or written by another person, or substantial proportions of material which have been accepted for the published of any other published materials, except where due acknowledgement is made in the article. Any contribution made to the research by others, with whom author[s] have work, is explicitly acknowledged in the article.

## Conflict of Interest Statement

The author[s] declare that this article was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Copyright Statement

Copyright © Author(s). This article is published under the Creative Commons Attribution (CC BY 4.0) licence. Anyone may reproduce, distribute, translate and create derivative works of this article (for both commercial and non-commercial purposes), subject to full attribution to the original publication and authors. The full terms of this licence may be seen at <http://creativecommons.org/licenses/by/4.0/legalcode>

## EDITORIAL TEAM

### Editor in Chief

Dr. Hindarto, Universitas Muhammadiyah Sidoarjo, Indonesia

### Managing Editor

Mochammad Tanzil Multazam, Universitas Muhammadiyah Sidoarjo, Indonesia

### Editors

Fika Megawati, Universitas Muhammadiyah Sidoarjo, Indonesia

Mahardika Darmawan Kusuma Wardana, Universitas Muhammadiyah Sidoarjo, Indonesia

Wiwit Wahyu Wijayanti, Universitas Muhammadiyah Sidoarjo, Indonesia

Farkhod Abdurakhmonov, Silk Road International Tourism University, Uzbekistan

Bobur Sobirov, Samarkand Institute of Economics and Service, Uzbekistan

Evi Rinata, Universitas Muhammadiyah Sidoarjo, Indonesia

M Faisal Amir, Universitas Muhammadiyah Sidoarjo, Indonesia

Dr. Hana Catur Wahyuni, Universitas Muhammadiyah Sidoarjo, Indonesia


Complete list of editorial team ([link](#))

Complete list of indexing services for this journal ([link](#))

How to submit to this journal ([link](#))

**Article information**

**Check this article update (crossmark)**



**Check this article impact (\*)**



**Save this article to Mendeley**



(\*) Time for indexing process is various, depends on indexing database platform

## **Systematic analysis of the dominant types of entomophages in fruit orchards**

**B.A. Sulaymonov, IsashovaPhD@mail.ru, (0)**

*Andijan branch of Tashkent state agrarian university, Uzbekistan*

**U.A. Isashova, IsashovaPhD@mail.ru, (0)**

*Andijan branch of Tashkent state agrarian university, Uzbekistan*

**M.K. Rakhmanova, mrbobursobirov@gmail.com, (0)**

*, Uzbekistan*

**M.Q.Parpiyeva, mrbobursobirov@gmail.com, (0)**

*, Uzbekistan*

**U.Sh. Rasulov, mrbobursobirov@gmail.com, (0)**

*, Uzbekistan*

<sup>(1)</sup> Corresponding author

### **Abstract**

This article is focused on a systematic analysis of leaf-mining fly types that belongs to Agromyzidae family. Due to it, 2 types of leaf-mining flies were identified in agricultural plants and 1 type in weeds. Also, it was identified that *Liriomyza sativae* Blanchard type of leaf-mining flies that belong to 4 families are met in 12 different plants in conditions of the Andijan region. Besides that, a number of insecticides as indoxamecty, protect were used against these kinds of pests.

Published date: 2019-12-07 10:11:10

## Introduction.

Over the last years a new type of pest appeared (it was registered in 1999 in our Republic). Its harm to agricultural products is increasing much. It is a leaf-mining fly that harms different vegetables, melons and other plants and adopts other plants as well. It was named as cosmopolite insect by the scientists of the world. Because of convenient climatic condition of our region this pest is adopting different plants and is spreading fast. One of the main reasons for this is there are not much natural fighters against it in our country[1]. Nowadays, Agromyzidae family is a representative of the group of two winged that harm most in plant growing. More than 300 types of this family are registered in the world's fauna. Most types of this family are considered to be monophage and 16 types as polyphages. 250 types of this family harm agricultural plants of the world. 100 types of them were registered to be met in former Soviet Union [2].

*Liriomyza sativae* Blanch(Diptera, Agromyzidae)type of leaf-mining flies have been harming agricultural products more than the other types in our Republic for the last period. Due to Zlobin and Drugova's conceptions *Liriomyza bryoniae* and *L. Strigata* type of leaf-mining flies can be met in Uzbekistan [3].

In 25 generations of Agromyzidae family *Liriomyza* is secondly important. A number of observations are carried out on determining the types of leaf-mining flies in Andijan. According to investigations and observations several patterns of leaf-mining flies were gathered from the fields with vegetables. Their types were identified with the help of the cooperation of specialists from Zoology Institutions.

The following types of leaf-mining flies were determined in the patterns taken from the plants of solanum family in the investigations carried out in Andijan region:

*Liriomyza sativae* Blanchard

*Liriomyza cicerinae* Rond

*Liriomyza strigata* Meigen

As it was identified in the carried observations leaf-mining flies that harm the representatives of solanum family are spread widely as a dominant type. Due to it, a basic part of leaf-mining flies is composed of *L. sativae* type. Also, *L. cicerinae* type was also determined in this territory. *L.strigata* type is mainly observed to harm weeds.

## Main part

*Liriomyza sativae* Blanch and *Liriomyza cicerinae* Rond types of leaf-mining flies were observed to harm agricultural products in biocenosis of our region. *Liriomyza huidobrensis* Blanch and *Liriomyza bryoniae* Kalt., *Chromatomyia fuscata* Zett., *Opomyza phaseolus*., *Phitomyza ilicis* Kert. may be spread in our Republic. *Liriomyza trifolii* Burg. Type is registered as an object of outer quarantine.

**Table 1** Systematic analysis of leaf-mining flies (Agromyzidae) and and frequency of their appearance (Andijan region, 2016-2017)

| №  | Types                                 | Group   | Family      | Registration |
|----|---------------------------------------|---------|-------------|--------------|
| 1  | <i>Liriomyza sativae</i> Blanch       | Diptera | Agromyzidae | +            |
| 2. | <i>Liriomyza trifolii</i> Burg.       | Diptera | Agromyzidae | -            |
| 3  | <i>Liriomyza strigata</i> Meigen.     | Diptera | Agromyzidae | +            |
| 4  | <i>Liriomyza cicerinae</i> Rodani     | Diptera | Agromyzidae | +            |
| 5. | <i>Liriomyza huidobrensis</i> Blanch. | Diptera | Agromyzidae | .*           |
| 6. | <i>Liriomyza bryoniae</i> Kalt.       | Diptera | Agromyzidae | .*           |
| 7. | <i>Chromatomyia fuscata</i> Zett.     | Diptera | Agromyzidae | .*           |
| 8. | <i>Opomyza phaseolus</i>              | Diptera | Agromyzidae | .*           |
| 9. | <i>Phitomyza ilicis</i> Kert.         | Diptera | Agromyzidae | .*           |

**Table 1.**

We continued our investigation in order to identify the frequency of the appearance of *L.satviae* type in various agricultural plants of Andijan region. The squares with the pests were identified and the types of plants were studied during 2016-2017.

It is seen from the following table, it is seen that when leaf-mining flies (*L.sativa*) are spread in a number of plants grown in Andijan, 4 families of leaf-mining flies harm 12 types of plants in different degrees[4].

**Table 2** Degree of harming the plant types by leaf-mining flies (*L.sativa*) (Andijan region, Andijan district, 2016-2017)

| №  | Type of a plant | Latin name of a plant       | Plant families | Degree of appearance |
|----|-----------------|-----------------------------|----------------|----------------------|
| 1  | Potato          | <i>Solanum lycopersicum</i> | Solanaceae     | +++                  |
| 2  | Egg-palnt       | <i>Solanum melongena</i>    | Solanaceae     | ++                   |
| 3  | Potato          | <i>Solanum tuberosum</i>    | Solanaceae     | ++                   |
| 4  | Pepper          | <i>Capsicum</i>             | Solanaceae     | ++                   |
| 5  | Bell-pepper     | <i>Capsicum annuum</i>      | Solanaceae     | ++                   |
| 6  | Peas            | <i>Cicer arietinum</i>      | Fabaceae       | +++                  |
| 7  | Water melon     | <i>Citrullus lanatus</i>    | Cucurbitaceae  | +++                  |
| 8  | Melon           | <i>Cucumis melo</i>         | Cucurbitaceae  | +++                  |
| 9  | Cucumber        | <i>Cucumis sativus</i>      | Cucurbitaceae  | +++                  |
| 10 | Pumpkin         | <i>Cucurbita maxima</i>     | Cucurbitaceae  | +++                  |
| 11 | Tobacco         | <i>Nicotiana tabacum</i>    | Solanaceae     | ++                   |
| 12 | Basilica        | <i>Ocimum basilicum</i>     | Lamiaceae      | +                    |

**Table 2.**

Comment: +++- much, ++-average, + little

4 representatives of Cucurbitaceae family are met in ( ), water melon ( ), cucumber ( ) and pumpkin plants ( and a representative of Solanaceae family is met in tomato( ), a representative of Fabaceae family is met in peas ( ) in high degree. Leaf-mining flies are met in average degree in egg plants of Solanaceae family ( ), bitter pepper ( ) and tobacco. They can be rarely met in the plants of Laminaceae family during the season.

In conclusion we can say that, 12 plants of solanum family (Solanaceae), pumpkins (Cucurbitaceae), leguminous plants (Fabaceae) are harmed with leaf-mining flies. Tomatoes, cucumbers and pumpkins are harmed much by these pests.

More than 330 types of *Liriomyza* generation are phytophages in agricultural plants all over the world and sometimes destroy the plants up to 75-80%. The most important economical harm is made in vegetable, leguminous crops and decorative gardening. In the practice of production chemicals are widely used to decrease the amount of pests in a short period of time. But some types of pests have high tolerance to insecticides. The researchers of different countries are carrying out a number of investigations in harmonized fighting against pests [5].

According to B.Sulaymonov's investigations, 3 classes, 12 groups, 12 families and 36 types of pests harm tomatoes and cucumbers in green houses of Uzbekistan. Here the harms of leaf-mining flies are very high. Such preparations as Sumi alfa 20% in 0,6l/hect solution, Konfidor 20% in 0,25 l/hect solution, Vermitek 1,8% in 0,6l/hect solution, Carbaphos 50% in 0,6l/hect solution show high effectiveness in chemical fighting against pests.[6].

Biology of *Liriomyza sativa* type was studied in Beijing region of Chinese People's Republic. Due to the results, the process of the lives of male and female pests lasted up to 12-20 days under 28 °C and 70% of relative air moist. They paired on the 2nd day of imago phase. Also, a number of preparations were tested. Xia yu and Ai Fu Ding preparations effected on worms and imagos in 80%-95% of cases. Ai Fu Ding preparation effected on chrysalis in 34% of cases [7]

Object and methods of investigation. We carried out investigations on chemical fighting against pests in the station of the institute of scientific investigations "Vegetable, melon and potato growing" in Andijan's scientific experimental station. In this experiment we used such chemical preparations as Indoxemactin (Indoxacarb 7.5%+Abamectin1.8%)0.2l/hect and Protect (Emmamectin benzoate 5%) 0.5kg/hect against leaf-mining flies. Vermitek preparation was taken as a pattern.

Our experiments were held in 4 versions and 4 repetitions. Before using the preparations, we identified the larva in 3 leaves of each plant. For this we took patterns of 20 different parts of each version. By this way we determined the number of larva in each leaf.



## Results of the investigation.

In the 1st version of our experiment we identified the existence of average 4.1 larva of pests in a leaf when Indoxamectin (Indoxacarb 7.5%+Abamectin 1.8%) preparations were used in the solution of 0.2 l/hect. On the 3rd day of our observations after the usage of the preparation the number of pests decreased up to 1.6. In this case the biologic effectiveness of chemical preparations was equal to 60.9%. On the 7th day of the experiment the number of pests in a leaf decreased up to 0.5 and the biologic effectiveness of the preparation composed 88.4%. On the 14th day of the experiment the influence of the chemical preparation a bit weakened. That is, the number of pests on the leaf a bit increased than on the 7th day. The number of pests on 1 leaf composed 0.9. The biologic effectiveness composed 79.7%. On the 21st day of the experiment the number of pests on 1 leaf was 1.3, biologic effectiveness composed 71.4%. On the 28th day of our observations the number of pests in 1 leaf was 2.0 and the biologic effectiveness was 57.1 %.

In the 2nd version where Protect (Emamectin benzoate 5%) preparation was used in 0,5 kg/hect solution, 4,2 pest larvae were identified in 1 leaf. On the 3rd day of our observations the number of pests decreased up to 1.9. In this case the biologic effectiveness of the preparation was 55.8%. On the 7th day of the experiment the number of pests in 1 leaf decreased to 0.7 and the biologic effectiveness of the preparation composed 85.8%. on the 14th day of the experiment the influence of the preparation a bit weakened.

Biologic effectiveness of chemicals against leaf-mining flies (*Liriomyza sativae*).

That is the number of pests in 1 leaf a bit increased than on the 7th day. The number of pests in 1 leaf composed 1.2 and the biologic effectiveness composed 76.7%. On the 21st day of the experiment it was identified that the number of pests in 1 leaf composed 1.7 and the biologic effectiveness composed 69.6%. On the 28th day of our observations, the number of pests on 1 leaf was 2.5 and the biologic effectiveness composed 55.3%.

In the 3rd version of our experiment where Vermitek 1.8% (Abamectin) preparation was used in 0.25 l/hect solution we identified the existence of average 4,5 larva of pests. On the 3rd day of our observations after the usage of preparation the number of pests decreased to 2.0. Here the biologic effectiveness of the chemical preparation composed 56.6 %. On the 7th day of the experiment the number of pests in 1 leaf decreased up to 0,9 and the biologic effectiveness of the preparation composed 83.0%. On the 14th day of the experiment it was identified that the influence of the chemical preparation a bit weakened. That is the number of pests in 1 leaf increased a bit than the 7th day of the experiment. The number of pests in 1 leaf composed 2.17 and the biologic effectiveness composed 69.3%. On the 21st day of the experiment the number of pests in 1 leaf composed 2.9, the biologic effectiveness composed 50.7%. On the 28th day of our observations the number of pests in 1 leaf was 3.4 and the biologic effectiveness composed 43.7%.

## Conclusion.

In our investigations we identified that *L. sativae* Blanch and *L. cicernae* Rond types of leaf-mining flies harm agricultural plants in agrobiocenosis of vegetables of Andijan region. When comparing the development of different types of leaf-mining flies in various plants, 4 representatives of Cucurbitaceae family harm melon, water melon, cucumber and pumpkin plants and a representative of Solanaceae family harm tomato much. In chemical fight against this pest Indoxamectin (Indoxacarb 7.5%+Abamectin 1.8%) was used in 0.2l/hect. On the 7th day of the experiment the number of pests in 1 leaf decreased to 0.7 and the biologic effectiveness of the preparation composed 88.4%. When Protect Emamectin benzoate 5%) was used in the solution of 0.5kg/hect. The number of pests decreased to 1.1 and the biologic effectiveness of the preparation composed 85.8%.

## References

1. Isashova U. - "Dangerous economical harm of leaf-mining flies in tomato plants of green houses (*Liriomyza sativae* Blanch)" *Agroscience*, 2-2018. -p 57.
2. Zlobin V.V., Drugova E.V. "Leaf-mining flies of Agromyzidae family". Moscow, Protection and quarantine of plants, 2002. № 7, -p 28.
3. Zlobin V.V., " To fauna of leaf-mining flies of Agromyzidae (Diptera) family in Zeys state reservation". 1986 . -p 51-53.
4. Isashova U. - "Damage of different types plant from percent of minimum flies (*Liriomyza sativae* Blanch)" *Актуальные проблемы современной науки*, № 4, 2018 Том 1.стр 212-215
5. Sujay, Y.H., Sattagi, H.N.,Patil, R.K., (2010), Invasive aliens insects and their impact on agro ecosystem, *Karnataka Journal of Agricultural sciences*, 23: 26-34
6. Sulaymonov B.A. "Pests of vegetable plants of green houses and management of their number.// *Agro ilm.*- Tashkent 2008.- №4(8). -p 26-28.
7. Wang Zin Zhong «Biology of *Liriomyza sativae* Blanchard and chemical control» *journal of Beijing agricultural college* 2003-04. -P. 189